

PERSONAL DATA

First Name: MICHELE
Surname: RIMOLDI
Born: CANTÙ (CO), 17/11/1984
Nationality: ITALIAN
Contact Address: DIPARTIMENTO DI SCIENZE MATEMATICHE
Politecnico di Torino
Corso Duca degli Abruzzi, 24
10129 Torino, ITALY
e-mail: WORK michele.rimoldi@polito.it
 PERSONAL michele.rimoldi@gmail.com
Personal page: <http://michelerimoldi.altervista.org>

CURRENT POSITION

Since January 2017 Researcher (RTD-a)
 Politecnico di Torino
 Dipartimento di Scienze Matematiche "Giuseppe Luigi Lagrange"

PREVIOUS POSITIONS

Oct. 2016-Dec. 2016 "Emma e Giovanni Sansone" Junior visiting position
 Centro di Ricerca Matematica "Ennio De Giorgi"
 Scuola Normale Superiore (Pisa, Italy)
Dec. 2015-Sept. 2016 Post-Doc fellowship (1 year)
 of the Fondation Sciences Mathématiques de Paris (FSMP)
 Laboratoire Analyse Géométrie et Applications (LAGA)
 Institut Galilée
 Université Paris 13 - Sorbonne Paris Cité
Jan. 2014 - Nov. 2015 Assegno di Ricerca (2 years Post-Doc position),
 Università degli Studi di Milano-Bicocca
 Dipartimento di Matematica e Applicazioni
Dec. 2011 - Dec. 2013 Assegno di Ricerca (2 years Post-Doc position),
 under the supervision of Prof. Stefano Pigola,
 Università degli Studi dell'Insubria (Como, Italy)
 Dipartimento di Scienza e Alta Tecnologia

ACADEMIC HABILITATIONS

December 2014 National Italian habilitation as *Professore di II fascia*
 in Geometry and Algebra. Sector: 01/A2.
 Validity: 30/12/2014–30/12/2020
February 2016 Qualification Française aux fonctions de *Maître de conférences*.
 25ème section.

March 2018 National Italian habilitation as *Professore di II fascia*
in Geometry and Algebra. Sector: 01/A2.
Validity: 30/03/2018–30/03/2024

EDUCATION

March 2012 Ph.D. in Mathematics,
Università degli Studi di Milano, Italy
Thesis–title: *Rigidity results for*
Lichnerowicz Bakry–Émery Ricci tensors
Supervisor: Prof. Stefano Pigola

September 2008 Laurea specialistica (master degree) cum laude in Mathematics
Università degli Studi dell’Insubria (Como, Italy)
Thesis–title: *Caratterizzazione di modelli mediante*
equazioni differenziali
Supervisor: Prof. Stefano Pigola

January 2007 Laurea triennale (bachelor) cum laude in Mathematics
Università degli Studi dell’Insubria (Como, Italy)
Thesis–title: *Costruzione di superfici minime complete*
tra due piani paralleli in \mathbb{R}^3
Supervisor: Prof. Stefano Pigola

July 2003 Diploma di Maturità
Liceo Scientifico Galileo Galilei (Erba (CO), Italy)

PUBLICATIONS

- [IRV] D. IMPERA, M. RIMOLDI, G. VERONELLI
Density problems for second order Sobolev spaces and cut-off functions
on manifolds with unbounded geometry
To appear on INTERNATIONAL MATHEMATICS RESEARCH NOTICES
(Preliminary version on [arXiv:1810.02578](https://arxiv.org/abs/1810.02578))
- [RV2] M. RIMOLDI, G. VERONELLI
Extremals of Log Sobolev inequality on non-compact manifolds and Ricci soliton structures
CALCULUS OF VARIATIONS AND PARTIAL DIFFERENTIAL EQUATIONS.
(2019) 58:66, doi: 10.1007/s00526-019-1518-z
- [IR3] D. IMPERA, M. RIMOLDI
Quantitative index bounds for translators via topology
MATHEMATISCHE ZEITSCHRIFT.
292 (2019), no.1-2, 513-527, doi: 10.1007/s00209-019-02276-y
- [IRS] D. IMPERA, M. RIMOLDI, A. SAVO
Index and first Betti number of f -minimal hypersurfaces and self-shrinkers
To appear on REVISTA MATEMATICA IBEROAMERICANA
(Preliminary version on [arXiv:1803.08268](https://arxiv.org/abs/1803.08268))
- [MMR] C. MANTEGAZZA, S. MONGODI, M. RIMOLDI
The Cotton tensor and the Ricci flow
GEOMETRIC FLOWS.
2 (2017), no. 1, 49-71, doi: 10.1515/geofl-2017-0001

- [IR2] D. IMPERA, M. RIMOLDI
Rigidity results and topology at infinity of translating solitons of the mean curvature flow
COMMUNICATIONS IN CONTEMPORARY MATHEMATICS.
19, 1750002 (2017), no. 6, [21 pages], doi: 10.1142/S021919971750002X
- [IR1] D. IMPERA, M. RIMOLDI
Stability properties and topology at infinity of f -minimal hypersurfaces
GEOMETRIAE DEDICATA
178 (2015), no. 1, 21-47, doi: 10.1007/s10711-014-9999-6
- [RV1] M. RIMOLDI, G. VERONELLI
Topology of steady and expanding gradient Ricci solitons via f -harmonic maps
DIFFERENTIAL GEOMETRY AND ITS APPLICATIONS
31 (2013), no. 5, 623-638, doi: 10.1016/j.difgeo.2013.06.001.
- [PR2] S. PIGOLA, M. RIMOLDI
Complete self-shrinkers confined into some regions of the space
ANNALS OF GLOBAL ANALYSIS AND GEOMETRY
45 (2014), no. 1, 47-65, doi: 10.1007/s10455-013-9387-8
- [MR] P. MASTROLIA, M. RIMOLDI
Some triviality results for quasi-Einstein manifolds and Einstein warped products
GEOMETRIAE DEDICATA
169 (2014), no. 1, 225-237, doi: 10.1007/s10711-013-9852-3
- [R2] M. RIMOLDI
On a classification theorem for self-shrinkers
PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY
142 (2014), no. 10, 3605-3613, doi: 10.1090/S0002-9939-2014-12074-0
- [MRR] P. MASTROLIA, M. RIGOLI, M. RIMOLDI
Some geometric analysis on generic Ricci solitons
COMMUNICATIONS IN CONTEMPORARY MATHEMATICS
15 (2013), no. 3, 1250058, 25 pp., doi: 10.1142/S0219199712500587.
- [CMMR] G. CATINO, C. MANTEGAZZA, L. MAZZIERI, M. RIMOLDI
Locally conformally flat quasi-Einstein manifolds
JOURNAL FÜR DIE REINE UND ANGEWANDTE MATHEMATIK
675 (2013), 181-189, doi: 10.1515/crelle.2011.183.
- [R1] M. RIMOLDI
A remark on Einstein warped products
PACIFIC JOURNAL OF MATHEMATICS
252 (2011), no. 1, 207-218, doi: 10.2140/pjm.2011.252.207
- [MRV] P. MASTROLIA, M. RIMOLDI, G. VERONELLI
Myers-type theorems and some related oscillation results
JOURNAL OF GEOMETRIC ANALYSIS
22 (2012), no. 3, 763-779, doi:10.1007/s12220-011-9213-0
- [PRRS] S. PIGOLA, M. RIGOLI, M. RIMOLDI, A. G. SETTI
Ricci almost solitons
ANNALI DELLA SCUOLA NORMALE SUPERIORE DI PISA. CLASSE DI SCIENZE
X (2011), no. 4, 757-799, doi: 10.2422/2036-2145.2011.4.01
- [PRiS] S. PIGOLA, M. RIMOLDI, A. G. SETTI
Remarks on non-compact gradient Ricci solitons
MATHEMATISCHE ZEITSCHRIFT
268 (2011), no. 3-4, 777-790, doi: 10.1007/s00209-010-0695-4

- [PR1] S. PIGOLA, M. RIMOLDI
Characterizations of model manifolds by means of certain differential systems
 CANADIAN MATHEMATICAL BULLETIN
 55 (2012), 632–645, doi:10.4153/CMB-2011-134-0

PREPRINTS

- [IR4] D. IMPERA, M. RIMOLDI
Index and first Betti number of f -minimal hypersurfaces: general ambients
 (submitted, [arXiv:1811.04009](#))
- [IPR] D. IMPERA, S. PIGOLA, M. RIMOLDI
The Frankel property for self-shrinkers from the viewpoint of elliptic PDE's
 (submitted, [arXiv:1803.02332](#))

TALKS, SEMINARS AND POSTERS

- September 2018 UNIVERSITÀ DI CAGLIARI, CAGLIARI, ITALY
Differential Geometry Workshop 2018
Harmonic maps, biharmonic maps, harmonic morphism and related topics
 Invited talk:
 “*The Frankel property for self-shrinkers from the view point of elliptic PDE's*”
- September 2018 UNIVERSITÄT HAMBURG, HAMBURG, GERMANY
Ricci flow, mean curvature flow and related singular flows
 Invited talk:
 “*The Frankel property for self-shrinkers from the view point of elliptic PDE's*”
- April 2017 UNIVERSITÀ DI TORINO, TORINO, ITALY
Differential Geometry Days
 Invited talk:
 “*Extremals of Log Sobolev inequalities on non-compact manifolds and Ricci soliton structures*”
- February 2017 SCUOLA NORMALE SUPERIORE, PISA, ITALY
Workshop su varietà reali e complesse
geometria, topologia e analisi armonica
 Invited talk:
 “*Extremals of Log Sobolev inequalities on non-compact manifolds and Ricci soliton structures*”
- December 2016 SCUOLA NORMALE SUPERIORE, PISA, ITALY
Seminari di Analisi Geometrica
 Seminar:
 “*Extremals of Log Sobolev inequalities on non-compact manifolds and Ricci soliton structures*”
- September 2016 UNIVERSITÀ DI CAGLIARI, CAGLIARI, ITALY
Complex and Riemannian Geometry days
 Invited talk:
 “*Extremals of Log Sobolev inequalities on non-compact manifolds and Ricci soliton structures*”
- June 2016 UNIVERSITÉ PARIS 13, PARIS, FRANCE
Groupe de travail Analyse semi-classique
 Seminar:

- “Stability properties and topology at infinity of f -minimal hypersurfaces”*
- April 2016 UNIVERSITÉ PARIS 7, PARIS, FRANCE
Séminaire de Géométrie
 Invited talk:
“Properness and boundedness properties of complete self-shrinkers for the mean curvature flow”
- Mars 2016 UNIVERSITÉ PARIS 13, PARIS, FRANCE
Séminaire EDP non linéaires
 Invited talk:
“Obstructions on the topology at infinity of translating solitons of the mean curvature flow”
- April 2015 UNIVERSIDAD DE GRANADA, GRANADA, SPAIN
Workshop on Geometric Flows
 Invited talk:
“Rigidity results and topology at infinity of translating solitons of the mean curvature flow”
- February 2015 UNIVERSITÀ DI CAGLIARI, CAGLIARI, ITALY
Seminari di Matematica
 Seminar:
“Complete self-shrinkers confined into some regions of the space”
- November 2014 UNIVERSITÀ DEGLI STUDI DI MILANO, MILANO, ITALY
PDE’s and Global Analysis @ UniMi
 Invited talk:
“Complete self-shrinkers confined into some regions of the space”
- October 2014 LEVICO TERME, TRENTO, ITALY
Progressi Recenti in Geometria Reale e Complessa - IX
 Invited talk:
“Complete self-shrinkers confined into some regions of the space”
- February 2014 UNIVERSIDADE FEDERAL DO CEARÀ, FORTALEZA, BRASIL
VII Workshop on Geometric Analysis
 Invited talk:
“Complete self-shrinkers confined into some regions of the space”
- April 2011 CENTRO DI RICERCA MATEMATICA ENNIO DE GIORGI, PISA, ITALY
Ricci Solitons Days in Pisa 2011
 Contributed talk:
“Triviality results for quasi-Einstein metrics and Einstein warped products”
- February 2011 UNIVERSIDAD DE GRANADA, SPAIN
Spanish-Japanese Workshop on Differential Geometry in Granada
 Poster:
“Triviality results for quasi-Einstein metrics and Einstein warped products”
- January 2011 UNIVERSITÀ DEGLI STUDI DELL’INSUBRIA, ITALY
Seminari di Analisi Geometrica
 Seminar:
“Triviality results for quasi-Einstein metrics and Einstein warped products”
- January 2011 UNIVERSITÀ DEGLI STUDI DELL’INSUBRIA, ITALY
Seminari di Analisi Geometrica
 Seminar:
“Quasi-Einstein metrics and Einstein warped products”

RESEARCH VISITS

| | |
|-----------------|---|
| February 2015 | UNIVERSITÀ DI CAGLIARI, CAGLIARI, ITALY (1 WEEK) Invited by Prof. S. Montaldo. |
| February 2014 | UNIVERSIDADE FEDERAL DO CEARÀ, FORTALEZA, BRASIL (2 WEEKS) Invited by Prof. G. P. Bessa. |
| Sept.-Dec. 2012 | INSTITUT HENRI POINCARÉ, PARIS, FRANCE (3 MONTHS) Guest for the research trimester in <i>Conformal and Kähler Geometry</i> Organizers of the trimester: Prof. M. Gursky, Prof. H. Hebey, Prof. F. Pacard, Prof. J. Viaclovsky. |
| January 2011 | SCUOLA NORMALE SUPERIORE, PISA, ITALY (1 WEEK) Invited by Prof. C. Mantegazza. |

TEACHING EXPERIENCE

| | | |
|-----------|---|---------------|
| 2006-2007 | Università degli Studi dell'Insubria Tutoring (16 hours) for the following courses: "Calcolo II", "Geometria I", "Algebra I" | COMO, ITALY |
| 2011 | Università degli Studi dell'Insubria Exercise classes (20 hours) for the course "Analisi Matematica II". Main lecturer: Prof. Alberto G. Setti | COMO, ITALY |
| 2012 | Università degli Studi dell'Insubria Exercise classes (10 hours) for the course "Geometria II". Main lecturer: Prof. Stefano Pigola | COMO, ITALY |
| 2013 | Università degli Studi dell'Insubria Exercise classes (16 hours) for the course "Geometria II". Main lecturer: Prof Stefano Pigola | COMO, ITALY |
| 2015 | Università degli Studi di Milano-Bicocca Exercise classes (24 hours) for the course "Geometria I". Main lecturer: Prof. Davide L. Ferrario | MILANO, ITALY |
| 2017 | Politecnico di Torino Exercise classes (20 hours) for the course "Algebra lineare e geometria". Main lecturer: Prof. C. Cumino | TORINO, ITALY |
| 2017 | Politecnico di Torino Exercise classes (20 hours) for the course "Algebra lineare e geometria". Main lecturer: Prof. M. Ferrarotti | TORINO, ITALY |
| 2017 | Politecnico di Torino Exercise classes (20 hours) for the course "Linear algebra and geometry". Language: English. Main lecturer: Prof. E. Carlini | TORINO, ITALY |
| 2017 | Politecnico di Torino Exercise classes (20 hours) for the course "Geometria differenziale e computazionale". Main lecturer: Prof. E. Musso | TORINO, ITALY |

| | | |
|------|---|---------------|
| 2018 | Politecnico di Torino Exercise classes (20 hours) for the course “Linear algebra and geometry”. Language: English. Main lecturer: Prof. E. Carlini | TORINO, ITALY |
| 2019 | Politecnico di Torino Exercise classes (40 hours) for the course “Algebra lineare e geometria”. Main lecturer: Prof. E. Musso | TORINO, ITALY |

FUNDING INFORMATION

| | |
|-----------|--|
| 2019- | Participant to the PRIN Project 2017 REAL AND COMPLEX MANIFOLDS: TOPOLOGY, GEOMETRY AND HOLOMORPHIC DYNAMICS Scientific coordinator: Prof. Filippo Bracci Coordinator of the local unity: Prof. Emilio Musso |
| 2018 | Recipient of the FFABR: ricercatori FONDO PER IL FINANZIAMENTO ATTIVITÀ BASE DI RICERCA FUNDED BY THE MIUR GRANT VALUE: 3.000 EU |
| 2017-2020 | Responsible for the STARTING GRANT PER GIOVANI RICERCATORI A TD A E B - I TRANCHE 2017 - RIMOLDI (53-RIL17RIMMIC) FUNDED BY THE POLITECNICO DI TORINO GRANT VALUE: 15.000 EU |
| 2014-2015 | Participant to the PRIN Project 2010-2011: VARIETÀ REALI E COMPLESSE: GEOMETRIA, TOPOLOGIA E ANALISI ARMONICA Scientific coordinator: Prof. Fulvio Ricci Coordinator of the local unity: Prof. Stefano Meda |
| 2014 | Participant to the INdAM-GNAMPA Project : PROPRIETÀ ANALITICHE E SPETTRALI DI VARIETÀ PESATE E APPLICAZIONI Coordinator: Debora Impera |
| 2013 | Coordinator of the INdAM-GNAMPA Project: FUNZIONI E MAPPE ARMONICHE: MISURA QUANTITATIVA DELL’INSIEME CRITICO, REGOLARITÀ E PROBLEMI DI DIRICHLET GRANT VALUE: 2.500 EU |
| 2010 | Participant to the INdAM-GNAMPA Project: TEORIA DEL POTENZIALE NON LINEARE SU VARIETÀ RIEMANNIANE E APPLICAZIONI GEOMETRICHE Coordinator: Prof. Stefano Pigola |

SOCIETY MEMBERSHIPS

| | |
|------------|------------------------|
| 2010-2018 | Member of INdAM-GNAMPA |
| Since 2019 | Member of INdAM-GNSAGA |

ORGANIZATION OF SCIENTIFIC EVENTS

- 2019 Member of the organizing committee of the conference:
 GEOMETRIC ANALYSIS, SUBMANIFOLDS
 AND GEOMETRY OF PDE'S
 To be held at Politecnico di Torino, Italy, September, 2019
 Scientific Committee: C. Arezzo, G. Besson, G. Manno, E. Musso,
 S. Salamon, C. Terng.
- 2019 Member of the organizing committee of the school:
 GEOMETRIC ANALYSIS ON RIEMANNIAN AND SINGULAR
 METRIC MEASURE SPACES
 To be held in Como, Italy, July 1-5, 2019
 Mini-courses by: A. Carlotto, A. Fraser, C. de Lellis, A. Ros, S. Wenger.
 Scientific Committee: G. Besson, S. Pigola, A. Setti, M. Troyanov.
- 2018 Member of the organizing committee of the mini-school/workshop:
 PLURIPOTENTIAL THEORY, GEOMETRIC ANALYSIS
 AND CALIBRATED GEOMETRY
 Università di Torino, Italy, May 2-4, 2018
 Mini-courses by: J Lotay, L. Mari.
- 2016 Member of the organizing committee of the school:
 GEOMETRIC ANALYSIS ON RIEMANNIAN AND SINGULAR
 METRIC MEASURE SPACES
 Como, Italy, July 11-15, 2016
 Mini-courses by: H.-D. Cao, N. Gigli, T. Ilmanen, R. Schoen, C. Sormani,
 X.-P. Zhu.
 Scientific Committee: G. Besson, S. Pigola, A. Setti, M. Troyanov.
- 2015 Member of the organizing committee of the workshop:
 A GEOMETRY DAY IN COMO 2015
 Como, Italy, Jan 9, 2015
 Talks by: Z. Djadli, A. Malchiodi, L. Mari, B. Nelli,
 R. Paoletti.
- 2014 Member of the organizing committee of the workshop:
 A GEOMETRY DAY IN COMO 2014
 Como, Italy, Jan 10, 2014
 Talks by: F. Bonsante, F. Coda-Marques, I. Holopainen, N. Gigli,
 C. Mantegazza, M. Rigoli.
- 2013 Member of the organizing committee of the school:
 GEOMETRIC ANALYSIS ON RIEMANNIAN AND SINGULAR METRIC SPACES
 Como, Italy, Sept 30-Oct 5, 2013
 Mini-courses by: S. Alexander, G. Carron, E. Hebey, U. Lang, A. Neves
 Scientific Committee: G. Besson, S. Pigola, A. Setti, M. Troyanov.

 CONFERENCES, WORKSHOPS AND SUMMER SCHOOLS ATTENDED

- January 2019 UNIVERSITÀ DEGLI STUDI DELL'INSUBRIA, COMO, ITALY
A Geometry Day in Como
- December 2018 UNIVERSITÀ DEGLI STUDI DI MILANO-BICOCCA, MILANO, ITALY
A day in Riemannian Geometry and related topics
- September 2018 UNIVERSITÀ DI CAGLIARI, CAGLIARI, ITALY
Differential Geometry Workshop 2018
Harmonic maps, biharmonic maps, harmonic morphism and related topics
- September 2018 UNIVERSITÄT HAMBURG, HAMBURG, GERMANY
Ricci flow, mean curvature flow and related singular flows

- May 2018 UNIVERSITÀ DI TORINO, TORINO, ITALY
Pluripotential Theory, Geometric Analysis and Calibrated Geometry
- February 2018 SCUOLA NORMALE SUPERIORE, PISA, ITALY
*Workshop su varietà reali e complesse:
geometria, topologia e analisi armonica*
- April 2017 UNIVERSITÀ DI TORINO, TORINO, ITALY
Differential Geometry Days
- February 2017 SCUOLA NORMALE SUPERIORE, PISA, ITALY
*Workshop su varietà reali e complesse:
geometria, topologia e analisi armonica*
- September 2016 UNIVERSITÀ DI CAGLIARI, CAGLIARI, ITALY
Complex and Riemannian Geometry Days
- July 2016 LAKE COMO SCHOOL FOR ADVANCED STUDIES, COMO, ITALY
*School in Geometric Analysis
Geometric Analysis on Riemannian and singular metric measure spaces*
- April 2015 UNIVERSIDAD DE GRANADA, GRANADA, SPAIN
Workshop on Geometric Flows
- March 2015 GOETHE UNIVERSITÄT, FRANKFURT AM MAIN, GERMANY
*Flow(ers) & Friends
A workshop on Geometric Analysis*
- January 2015 UNIVERSITÀ DEGLI STUDI DELL'INSUBRIA, ITALY
A geometry day in Como
- November 2014 UNIVERSITÀ DEGLI STUDI DI MILANO, MILANO, ITALY
PDE's and Global Analysis @ UniMi
- October 2014 LEVICO TERME (TN), ITALY
Progressi Recenti in Geometria Reale e Complessa - IX
- September 2014 VILLASIMIUS (CA), ITALY
New trends in Differential Geometry 2014
- June 2014 CENTRO DE GIORGI, PISA, ITALY
ERC School on Geometric Evolution Problems
- February 2014 UNIVERSIDADE FEDERAL DO CEARÀ, FORTALEZA, BRASIL
VII Workshop on Geometric Analysis
- January 2014 UNIVERSITÀ DEGLI STUDI DELL'INSUBRIA, ITALY
A geometry day in Como
- October 2013 INSTITUT FOURIER, GRENOBLE, FRANCE
*Autumn days in Grenoble
Curvature in metric spaces*
- October 2013 LAKE COMO SCHOOL FOR ADVANCED STUDIES, COMO, ITALY
*School in Geometric Analysis
Geometric Analysis on Riemannian and singular metric spaces*
- June 2013 UNIVERSIDAD DE GRANADA, SPAIN
Variational problems and Geometric PDE's

| | |
|-----------------|--|
| Feb.-Mar. 2013 | SCUOLA NORMALE SUPERIORE, PISA, ITALY <i>Workshop su varietà reali e complesse: geometria, topologia e analisi armonica</i> |
| January 2013 | UNIVERSITÀ DEGLI STUDI DELL'INSUBRIA, ITALY <i>A geometry day in Como</i> |
| Sept.-Dec. 2012 | INSTITUT HENRI POINCARÉ, PARIS, FRANCE <i>Research trimester in Conformal and Kähler geometry</i> |
| June 2012 | ICTP, TRIESTE, ITALY <i>ICTP-ESF School and Conference on Geometric Analysis</i> |
| September 2011 | UNIVERSIDAD DE GRANADA, SPAIN <i>VI International Meeting on Lorentzian Geometry</i> |
| May 2011 | UNIVERSITÀ DI MILANO BICOCCA, ITALY <i>Geometria in Bicocca</i> |
| April 2011 | CENTRO DE GIORGI, PISA, ITALY <i>Ricci Solitons Days in Pisa 2011</i> |
| February 2011 | UNIVERSITÀ DI MILANO BICOCCA, ITALY 1st Bicocca HART <i>Harmonic Analysis and Related Topics</i> |
| February 2011 | UNIVERSIDAD DE GRANADA, SPAIN <i>Spanish-Japanese Workshop on Differential Geometry</i> |
| September 2010 | VERBANIA, ITALY Riemann International School of Mathematics <i>Nonlinear Differential Equations</i> |
| June-July 2010 | CETRARO (CS), ITALY CIME course on <i>Ricci Flow and Geometric Applications</i> |
| May 2010 | DRESDEN UNIVERSITY OF TECHNOLOGY, GERMANY The 8th AIMS Conference on <i>Dynamical Systems, Differential Equations and Applications</i> |
| September 2009 | UNIVERSITÀ DI CAGLIARI, ITALY <i>A harmonic map fest</i> |
| June 2009 | CENTRO DE GIORGI, PISA, ITALY Research Trimester on <i>Geometric flows and Geometric Operators</i> |
| April 2009 | VERBANIA, ITALY Riemann International School of Mathematics <i>Advances in number theory and geometry</i> |

REFeree ACTIVITY

Referee for: Proceedings of the Edinburgh Mathematical Society, Communications in Analysis and Geometry, Journal of Geometric Analysis, Journal of Mathematical Analysis and Applications, Illinois Journal of Mathematics, Manuscripta Mathematica, International Journal of Mathematics, Bulletin des Sciences Mathématiques, Annali di Matematica Pura ed Applicata, Differential Geometry and its Applications, Annals of Global Analysis

and Geometry, Bulletin of the Belgian Mathematical Society, International Journal of Geometric Methods in Modern Physics, Mathematische Nachrichten, Mediterranean Journal of Mathematics, Publicationes Mathematicae Debrecen, Monatshefte für Mathematik, Beirträge zur Algebra und Geometrie.

Reviewer for AMS Mathematical Reviews.

OTHER ACTIVITIES

Febr-June 2017

ATTENDANCE OF THE COURSE

Apprendere ad insegnare nell'Higher Education

Commitment: 40 hours.

Politecnico di Torino

LANGUAGE SKILLS

Mother tongue: Italian.

Other languages: English (Cambridge First Certificate in English (FCE), 2002).

RESEARCH ACTIVITY

So far, my research activity has mainly dealt with Riemannian Geometry, Geometric Analysis, and Geometric PDE's. In particular, I've been focusing on the following topics:

- **Ricci solitons.** ([PRiS], [PRRS], [MRR], [RV1], [RV2]) Let $(M, \langle \cdot, \cdot \rangle)$ be a Riemannian manifold. A Ricci soliton structure on M is the choice of a smooth vector field X (if any) satisfying the soliton equation $Ric + \frac{1}{2}\mathcal{L}_X \langle \cdot, \cdot \rangle = \lambda \langle \cdot, \cdot \rangle$. In the special case where $X = \nabla f$ for some smooth function $f : M \rightarrow \mathbb{R}$ we say that $(M, \langle \cdot, \cdot \rangle, \nabla f)$ is a gradient Ricci soliton with potential f . In this case the soliton equation reads $Ric_f = Ric + Hess(f) = \lambda \langle \cdot, \cdot \rangle$. Since the appearance of the seminal works of R. Hamilton, [25], and G. Perelman, [34], the study of gradient Ricci solitons has become the subject of a rapidly increasing investigation mainly directed towards two goals, classification and triviality.

(i) In collaboration with Stefano Pigola and Alberto G. Setti, in [PRiS] we have studied triviality and rigidity results and curvature estimates for gradient Ricci solitons under weighted L^p conditions on the relevant quantities. These extend and generalize, often in a significant way, previous results (see e.g. [16, 35, 37, 49, 11]). Moreover, we show how techniques coming from stochastic analysis, such as stochastic completeness, in the form of the weak Omori–Yau maximum principle, parabolicity and L^p -Liouville type results, once translated into the setting of weighted geometry, are natural in the investigation of gradient Ricci solitons, and lead to elegant proofs of the above mentioned results.

(ii) In collaboration with Stefano Pigola, Marco Rigoli, and Alberto G. Setti, in [PRRS] we have introduced an extension of the concept of gradient Ricci soliton, the Ricci almost soliton, allowing λ in the soliton equation to be a generic smooth function on the weighted manifold $M_f = (M, \langle \cdot, \cdot \rangle, e^{-f} dvol)$. In view of the fact that the soliton function λ is not necessarily constant, one expects that a certain flexibility on the almost soliton structure is allowed and, consequently, the existence of almost solitons is easier to prove than in the classical situation. This feeling is confirmed by a number of different examples of almost solitons. On the other hand we prove a rigidity result which indicates that almost solitons should reveal a reasonably broad generalization of the fruitful concept of classical soliton. In particular, one obtains that not every complete manifold supports an almost soliton structure. The investigation in [PRRS] is mainly concerned with triviality, pointwise curvature estimates and isolation phenomena for complete gradient Ricci almost solitons and the results follow from considering elliptic equations or inequalities for various geometric quantities on almost solitons and rely on analytical techniques. This is the same philosophy used e.g. by M. Eminenti, G. La Nave and C. Mantegazza in [16] or in [PRiS]. More specifically, one sees that the differential (in)equalities at hand naturally involve the f -Laplacian. Since the almost soliton equation means precisely that the f -Bakry–Émery Ricci curvature Ric_f is proportional to the metric tensor, we are naturally led to introduce a number of weighted manifolds tools whose range of application goes beyond the investigation of almost solitons. An important instance of these tools is represented by the Omori–Yau maximum principle for the f -Laplacian under both weighted Ricci lower bounds and weighted volume growth conditions. Moreover we extended to almost solitons also some topological results known in the classical case ([46, 19, 31, 16]).

(iii) While gradient Ricci solitons were widely studied, relatively little is known about generic Ricci solitons, that is, when the soliton field is not necessarily the gradient of a potential, and the majority of the results is concerned with the compact case. A first important difference is that, in the general case, we cannot make use of the weighted manifold structure which naturally arises when dealing with gradient solitons. The same applies for related concepts such as the Bakry–Émery Ricci tensor, whose boundedness from below with a suitable radial function, together with an additional assumption on the potential function gives rise to weighted volume estimates. These facts restrict the applicability of analytical tools such as the weak Omori–Yau maximum principle for the diffusion operator known as f -Laplacian, weighted L^p -Liouville-type theorems and *a priori* estimate. Nevertheless, in the general case the soliton structure is encoded in the geometry of an appropriate operator Δ_X that we shall call the X -Laplacian and that is defined on $u \in C^2(M)$ by $\Delta_X u = \Delta u - \langle X, \nabla u \rangle$. Assuming a suitable growth condition on the vector field X , in collaboration with Paolo Mastrolia and Marco Rigoli, in [MRR] we obtained scalar curvature and volume estimates and a gap theorem for the traceless Ricci tensor. These results permit to deduce some interesting geometric consequences.

(iv) In collaboration with Giona Veronelli, in [RV1], we studied f -harmonic maps from non-compact manifolds into non-positively curved ones. Notably, we proved existence and vanishing results which generalize to the weighted setting part of Schoen and Yau's theory of harmonic maps, [40, 41]. As an application, we deduced information on the topology of manifolds with lower bounded Bakry–Émery Ricci tensor, and in particular of steady and expanding gradient Ricci solitons. Unlike most results so far known some topological theorem is obtained without requiring any assumption on the weight function f .

(v) In collaboration with Giona Veronelli, on [RV2], we established the existence of extremals for the Log Sobolev functional on complete non-compact manifolds with Ricci curvature bounded from below and strictly positive injectivity radius, under a condition near infinity. This generalizes a result previously obtained in [47] where it was assumed that the manifold had bounded geometry (i.e. the Riemann curvature tensor and all its covariant derivatives are bounded and that there is a uniform positive lower bound on the volume of geodesic balls of radius 1). When Ricci curvature is also bounded from above we get exponential decay at infinity of the extremals. As a consequence of these analytical results we establish, under the same assumptions, that non-trivial shrinking Ricci solitons support a gradient Ricci soliton structure. Our technique hence gives different conditions from those previously considered in [31] for the validity of a such a conclusion. On the way, we prove two results of independent interest: the existence of a distance-like function with uniformly controlled gradient and Hessian on complete non-compact manifolds with bounded Ricci curvature and strictly positive injectivity radius (this result was actually generalized in a subsequent work also to manifolds with possibly unbounded geometry; see below) and a general growth estimate for the norm of the soliton vector field on manifolds with bounded Ricci curvature.

- **Quasi-Einstein manifolds.**([R1], [MR], [CMMR]) Another possible generalization of Einstein manifolds and gradient Ricci solitons, first considered by J. Case, Y.-S. Shu and G. Wei, [9], are quasi-Einstein manifolds. These are weighted manifolds M_f such that their modified Bakry–Émery Ricci tensor $Ric + Hess(f) - \frac{1}{k} df \otimes df$ is a constant multiple of the metric. The importance of these geometric structures comes from a problem (proposed in [3]) on the existence of Einstein manifolds realized as warped products. Indeed, in [29], D.-S. Kim and Y.-H. Kim proved a characterization of quasi-Einstein metrics as base metrics of Einstein warped product metrics.

(i) Exploiting this relationship, in [R1] and in [MR] (this latter written in collaboration with Paolo Mastrolia), we have obtained triviality and rigidity results for non-compact quasi-Einstein manifolds and Einstein warped products with non-compact bases and Einstein fibre. The proofs rely on the Omori–Yau maximum principle for the weighted Laplacian, a new gradient estimate for solutions of weighted Poisson-type equations and some Liouville-type theorems in the setting of weighted manifolds.

(ii) We have next faced the problem of the local characterization of locally conformally flat quasi-Einstein manifold. In particular, in collaboration with Giovanni Catino, Carlo Mantegazza, and Lorenzo Mazziere, in [CMMR] we proved that any complete locally conformally flat quasi-Einstein manifold of dimension $m \geq 3$ is locally a warped product with $(m - 1)$ -dimensional fibers of constant curvature. Although the result was already known for gradient Ricci solitons (of which quasi-Einstein manifolds are a generalization), [48, 10, 7], the strategy of the proof is completely new and can be used as the main step to classify locally conformally flat shrinking and steady gradient Ricci solitons.

- **Self-shrinkers and translators of the mean curvature flow: classification and rigidity results; Frankel property.**([R2], [IR2],[PR2], [IPR]) By a self-shrinker of the MCF we mean a connected, isometrically immersed hypersurface $x : \Sigma^m \rightarrow \mathbb{R}^{m+1}$ whose mean curvature vector field \mathbf{H} satisfies the equation $x^\perp = -\mathbf{H}$. Self-shrinkers play an important role in the study of mean curvature flow, since they describe all possible blow ups at a given singularity of a MCF. Let $f = \frac{|x|^2}{2}$ and denote by \mathbf{A} the second fundamental form of the immersion. In [R2] we observed that by the definition and Gauss' equation one gets that

$Ric_f \geq 1 - |\mathbf{A}|^2$. This important estimate permits to apply, once again, the analytical tools developed so far in weighted geometry. Indeed the differential (in)equalities at hand on self-shrinkers involve the natural drifted Laplacian Δ_f on Σ_f .

(i) The classification of mean convex self-shrinkers began with [27], where G. Huisken showed that the only smooth closed mean convex self-shrinkers are round spheres. In the complete non-compact case G. Huisken, [26], also proved a classification theorem saying that the only possible smooth mean convex embedded self-shrinkers with bounded $|\mathbf{A}|$ and (extrinsic) polynomial volume growth are the cylindrical products $C_{\sqrt{k}}^{k,m-k} = S_{\sqrt{k}}^k \times \mathbb{R}^{m-k}$, $k = 0, \dots, m$. Recently T. Colding and W. Minicozzi, [14], showed that the hypothesis $|\mathbf{A}|$ bounded can be dropped. The hypothesis of polynomial volume growth in [14] is used to show that various weighted integrals converge in order to justify various integration by parts and it is natural in the study of the singularities that a MCF goes through since any time-slice of a blow up of a closed MCF has polynomial growth. Nevertheless, looking only at the self-shrinker equation, it might be thought under what weaker conditions the conclusion in [14] still holds. In the main result in [R2] we replace polynomial volume growth with a weighted L^2 -condition on $|\mathbf{A}|$. The viewpoint of weighted manifolds we adopt permits also to recover easily a classification result by H.D. Cao and H. Li, [7], where instead of the mean convexity condition it is considered a L^∞ -type condition on $|\mathbf{A}|$.

(ii) By a translator of the MCF we mean a connected isometrically immersed complete hypersurface $x : \Sigma^m \rightarrow \mathbb{R}^{m+1}$ whose mean curvature vector field \mathbf{H} satisfies the equation $\mathbf{H} = v^\perp$ for some fixed unit length vector $v \in \mathbb{R}^{m+1}$. These hypersurfaces correspond to translating solitons of the MCF, and play a key role in the study of slowly forming singularities. Translators of the MCF turn out to be f -minimal hypersurfaces (critical points of a weighted area functional) in the Euclidean space \mathbb{R}^{m+1} with the density $e^{\langle v, x \rangle}$. In collaboration with Debora Impera, in the first part of the paper [IR2], we highlighted how the realm of weighted manifolds and f -minimal hypersurfaces can naturally give strong enough characterization results also in this setting. For example, we proved a rigidity theorem for f -stable translators under a weighted L^2 -condition on the norm of the second fundamental form, thus improving some previous rigidity results; see [42].

(iii) It was conjectured by H. D. Cao, [8], that every complete self-shrinker has extrinsic polynomial volume growth. By a result due to Q. Ding and Y. L. Xin, [15], later completed by X. Cheng and D. Zhou, [13], this is equivalent to the fact that the immersion is proper. As a first consequence of this result, one obtains easily a counterexample to Cao conjecture just taking the product of one of the non-closed self-shrinking curves in Abresch-Langer analysis and an Euclidean factor. Another consequence is that if a complete self-shrinker has extrinsic polynomial growth and it is confined in a ball of \mathbb{R}^{m+1} , then it should be compact. In collaboration with Stefano Pigola, in [PR2] we moved some steps in getting information on the geometry of non-compact bounded self-shrinkers (if any) obtaining natural and general restrictions that force these hypersurfaces to be compact (e.g. global boundedness conditions, pinching conditions at infinity and $L^{p \geq m}$ conditions on $|\mathbf{A}|$). More generally we tried to understand the geometry of self-shrinkers which are confined in a connected region bounded by some dilated cylinder $C_R^{k,m-k}$. This was done proving a general strong maximum principle for self-shrinkers. Actually, when $k = 0$, it is reasonable that a complete self-shrinker has a certain homogeneous distribution around $0 \in \mathbb{R}^{m+1}$ and, therefore, it should intersect every hyperplane through the origin. Moreover, by the maximum principle, the intersection should be transversal somewhere unless the self-shrinker is itself an hyperplane. For compact self-shrinkers these properties are easily verified and a similar conclusion can be obtained for complete self-shrinkers with a controlled extrinsic geometry. In [PR2] we pointed out natural geometric conditions that permit to recover the full conclusion of the compact case. Moreover, once we have understood that, to a certain extent, complete self-shrinkers intersect transversally a hyperplane through the origin, we were also able to deduce spectral information on the drifted Laplacian Δ_f whenever the intersection is compact, and some (extrinsic) volume growth condition is satisfied. Some of the ideas introduced in this paper were later also applied in [IPR] to give a direct proof of the fact that a properly immersed self-shrinker cannot be located neither inside nor outside a self-shrinker cylinder.

(iv) In many instances, properly immersed self-shrinkers of the MCF behave like closed minimal hypersurfaces of the standard sphere. In this latter setting, it is well known that any two closed minimal immersed hypersurfaces must intersect. Actually the ambient space can be generalized to a compact Riemannian manifold with strictly positive Ricci curvature. This is called the Frankel property after the celebrated paper [20] by T. Frankel. Starting from the work by G. Wei and W. Wylie, [45], where the case of compact hypersurfaces is considered, in collaboration with D. Impera and S. Pigola in [IPR] we investigated the validity of the (smooth) properly embedded Frankel property for self-shrinkers. In particular, we proved that two properly embedded self-shrinkers in the Euclidean space that are sufficiently separated at infinity must intersect at a finite point. The proof argues by contradiction and is based on a localized version of the Reilly formula applied to a suitable f -harmonic function with controlled gradient which is shown to exist,

under our assumptions, in the hypothetical region which separates the two self-shrinkers.

- **f -minimal hypersurfaces, self-shrinkers and translators of the MCF: stability properties and their interactions with topology at infinity and topological invariants** ([IR1], [IR2], [IRS], [IR3], [IR4]) Following M. Gromov, [24], if we consider an isometrically immersed orientable hypersurface Σ^m in the weighted manifold M_f^{m+1} , we can define a generalization of the mean curvature vector field as $\mathbf{H}_f = \mathbf{H} + (\nabla f)^\perp$. Since the weighted structure on M induces a weighted structure on Σ we can consider the variational problem for the weighted area functional. From variational formulae, one can see that Σ is f -minimal, namely a critical point of the weighted area functional, if and only if \mathbf{H}_f vanishes identically. Clearly minimal hypersurfaces are a particular case of f -minimal hypersurfaces corresponding to the case $f \equiv \text{const}$. Moreover, self-shrinkers of the mean curvature flow are important examples of f -minimal hypersurfaces, when the ambient weighted space is the Euclidean space endowed with the Gaussian measure $e^{-\frac{|x|^2}{2}} d\text{vol}_{\mathbb{R}^{m+1}}$.

(i) The research on f -minimal hypersurfaces has started in recent years and it has been already approached by many authors, see e.g. [18], [12], [17]. Much efforts have been devoted to the study of the stability properties. These are taken into account by spectral properties of the weighted Jacobi operator $L_f = -\Delta_f - (|\mathbf{A}|^2 + \text{Ric}_f(\nu, \nu))$. The most up to date result, proved by X. Cheng, T. Mejia, and D. Zhou, [12], states that there exist no L_f -stable complete f -minimal hypersurfaces Σ immersed in a complete weighted Riemannian manifold M_f with $\text{Ric}_f \geq k > 0$, provided $\text{vol}_f(\Sigma) < +\infty$. In collaboration with Debora Impera, in [IR1], we generalized this result considering growth conditions on the intrinsic weighted volume of geodesic balls. Furthermore, in the instability case, exploiting the oscillatory behaviour of solutions of some ODEs that naturally arise in this setting, we investigated general geometric restrictions for the finiteness of the weighted index of the f -minimal hypersurface, that is, the maximum dimension of the linear space of compactly supported deformations that decrease the weighted area up to second order. Moreover, exploiting a weighted version of a finiteness result and the adaptation to this setting of Li-Tam theory, we investigated the topology at infinity of f -minimal hypersurfaces. On the way, we proved a new comparison result in weighted geometry and we provided a general weighted L^1 -Sobolev inequality for hypersurfaces in Cartan-Hadamard weighted manifolds satisfying suitable restrictions on the weight function and on its radial derivative.

(ii) In the particular case of translators of MCF, in [IR2], we obtained the validity of a weighted L^1 Sobolev inequality on translators which are contained in a upper halfspace with respect to the translating direction. The validity of this inequality permits to obtain very neat results on the topology at infinity of f -stable translators and translators with finite f -index. The main tools here are again weighted versions of Li-Tam theory and of an abstract finiteness results, which were developed in [IR1]. Moreover, as a consequence of the weighted L^1 -Sobolev inequality, the assumption on the finiteness of the f -index can be also replaced by the assumption that the hypersurface has finite weighted total curvature.

(iii) The results obtained in [IR2] have a qualitative nature. Inspired by the recent work by C. Li in the setting of minimal hypersurfaces in the Euclidean space, [30], in [IR3] we combined the weighted Li-Tam theory discussed above and a technique pioneered by A. Savo and A. Ros, [39], [38], to obtain a quantitative estimate on the generalized index of translator for the mean curvature flow with bounded norm of the second fundamental form. The estimate involves the dimension of the space of weighted square-integrable f -harmonic 1-forms, and thus yields estimates in terms of the number of ends of the hypersurface when this is contained in a upper halfspace with respect to the translating direction. When there exists a point where all principal curvatures are distinct we estimated the nullity of the stability operator. This permitted to obtain quantitative estimates on the stability index via the topology of translators with bounded norm of the second fundamental form which are either two-dimensional or (in higher dimension) have finite topological type and are contained in a upper halfspace.

(iv) The recent impressive developments in the existence theory for minimal immersions through min-max method have motivated a renewed interest in studying estimates on the Morse index of minimal immersions. One possible way to control instability is through topological invariants (in particular the first Betti number) of the minimal hypersurface. This was first investigated by A. Ros in [38] for immersed minimal surfaces in \mathbb{R}^3 , or a quotient of it by a group of translations, and then, in higher dimension, by A. Savo when the ambient manifold is the round sphere, [39]. The idea behind these works is to use harmonic 1-forms and some ambient structure to make some interesting functions out of those, permitting to get a lower bound on the index. In collaboration with Debora Impera and Alessandro Savo, in [IRS], we employed the method in [39] to study the Morse index of self-shrinkers for the mean curvature flow and, more generally, of f -minimal hypersurfaces in a weighted Euclidean space endowed with a convex weight. When the hypersurface is compact, we showed that the index is bounded from below by an affine function of its first Betti number. When the first Betti number is large this improves index estimates known in literature. In the complete non-compact case, the lower bound is in terms of the dimension of the space of weighted square integrable

f -harmonic 1-forms; in particular, in dimension 2, the procedure gives an index estimate in terms of the genus of the surface.

(v) The technique in [38] and [39] was recently extended and generalized in [2], where it is shown, from a general perspective that the Morse index is bounded from below by a linear function of the first Betti number for all closed minimal hypersurfaces in a large class of positively curved ambient manifolds. This comprises, for instance, all compact rank one symmetric spaces. In their work, the ambient structure is taken into account by the existence of some nice isometric immersion into an Euclidean space, via which one can produce instability directions, again using harmonic 1 forms. Following this idea and motivated by the approach introduced in [IRS], in [IR4] we studied quantitative Morse index estimates in the wider setting of closed f -minimal hypersurface sitting in a general weighted manifold. The technique permits, in particular, to obtain a linear lower bound on the Morse index via the first Betti number for closed f -minimal hypersurfaces in products of some compact rank one symmetric spaces with an Euclidean factor, endowed with the rigid shrinking gradient Ricci soliton structure. These include, as particular cases, all cylindrical shrinking gradient Ricci solitons.

- **Density problems for Sobolev spaces and second order distance-like on manifolds with unbounded geometry.** ([IRV]) Let (M, g) be a complete non-compact Riemannian manifold. The behaviour of the distance function from a fixed reference point $r(x)$ can reflect how different the manifold is from the Euclidean space. In general $r(x)$ is 1-Lipschitz on M , but only a.e. differentiable. However, a well-known result by Greene and Wu shows the existence of a function $H(x)$ which is smooth, distance-like (i.e. $r(x)/C < H(x) < Cr(x)$), and whose gradient is bounded. It is a natural question to what extent one could hope to generalize this result, giving sufficiently general geometric assumptions implying the existence of a distance-like function with controlled higher order derivatives. In collaboration with D. Impera and G. Veronelli, in [IRV] we considered complete non-compact manifolds which satisfy either a sub-quadratic growth of the norm of the Riemann curvature, or a sub-quadratic growth of both the norm of the Ricci curvature and the squared inverse of the injectivity radius. We show the existence on such a manifolds of a distance-like function with bounded gradient and mild growth of the Hessian. As a main application, we proved that smooth compactly supported functions are dense in $W^{2,p}$. The result is improved for $p = 2$ avoiding both the upper bound on the Ricci tensor, and the injectivity radius assumption. As further applications we prove new disturbed Sobolev and Calderón-Zygmund inequalities on manifolds with possibly unbounded curvature and highlight consequences about the validity of the full Omori-Yau maximum principle for the Hessian.
- **Myers' type theorems.** ([MRV]) We analyzed the roots of the solutions of the special ODE $u''(t) + k(t)u(t) = 0$. The existence of (at least) a zero for a solution of this differential equation implies the compactness of the m -dimensional manifold M , once $(m-1)k(t) = \text{Ric}_\gamma(t)$, where Ric_γ is the radial Ricci curvature of M . Developing an idea of E. Calabi, [6], in collaboration with Paolo Mastrolia and Giona Veronelli, in [MRV] we gave a new compactness result for complete manifolds with Ricci curvature lower bounded by a negative constant, thus extending previous results in [1, 21, 32]. Similar techniques permit to analyze also the zeros of the more general problem $(v(t)u'(t))' + k(t)v(t)u(t) = 0$, for some "weight" function $v(t)$. Even in this case, we studied assumptions which guarantee existence of a zero and oscillation of the solutions. As previously observed by B. Bianchini, L. Mari and M. Rigoli, [4], an application of these results yields new estimates on the bottom of the spectrum and on the index of some Schrödinger operator on Riemannian manifolds.
- **Evolution by Ricci flow of geometric tensors.** ([MMR]) Complete gradient steady Ricci solitons play an important role in the study of Hamilton's Ricci flow as they correspond to translating solutions, and often arise as type II singularity models. Thus one is interested in classifying them and understanding their geometry. R. Bryant has discovered a steady Ricci soliton in dimension 3, which is rotationally symmetric and has positive curvature operator. Bryant's construction can be adapted to higher dimensions. Inspired by the uniqueness property in dimension 2 of the cigar soliton, in [35], Perelman conjectured that the Bryant soliton is (up to scaling) the only complete non-compact three-dimensional gradient steady Ricci soliton with positive sectional curvature which satisfies a non-collapsing assumption at infinity (namely \mathcal{K} -noncollapsness). In [7]. H.D. Cao and Q. Chen proved uniqueness in dimension $m \geq 3$ under the additional assumption that (M^m, g) is locally conformally flat. The same result was proved independently by G. Catino and C. Mantegazza, [10], in dimension $m \geq 4$. Then, increasingly weaker conditions were considered, until, very recently, S. Brendle gave a proof of the conjecture without any further assumption, [5].

The proof strategy adopted in [10] to obtain uniqueness of locally conformally flat steady gradient non-flat Ricci solitons (for $m \geq 4$) is based on the computation of the evolution of the Weyl tensor under the Ricci flow. Inspired by this technique, in collaboration with C. Mantegazza and S. Mongodi, [CMM], we have

computed the evolution of the Cotton and the Bach tensor under the Ricci flow of a Riemannian manifold, with particular attention to the three-dimensional case and we discussed some applications. Beyond their own interest, these computations possibly suggest an alternative way to address Perelman's conjecture.

- **Obata's type characterization theorems.**([PR1]) We studied rigidity properties of Riemannian manifolds supporting solutions of particular second order differential systems. In particular, in collaboration with Stefano Pigola, in [PR1] we have considered more general differential systems than the classical ones of Obata's type ([33, 28, 44]), allowing the coefficient for the linear term to be a function (non-necessarily constant) of the distance function from a reference point. This permitted us to extend the characterization of space-forms to the case of model manifolds. We generalized also some metric rigidity results ([22]) by means of equations involving (gradient) vector fields.

Bibliography for the research activity

- [1] W. Ambrose, *A theorem of Myers*, Duke Math. J. **24** (1957), 345–348.
- [2] L. Ambrozio, A. Carlotto, B. Sharp, *Comparing the Morse index and the first Betti number of minimal hypersurfaces*, J. Differential Geom. **108** (2018), no. 3, 379–410.
- [3] A. Besse, *Einstein manifolds*, Springer–Verlag, Berlin–Heidelberg (1987).
- [4] B. Bianchini, L. Mari, M. Rigoli, *Spectral radius, index estimates for Schrödinger operators and geometric applications, (English summary)*, J. Funct. Anal. **256** (2009), no. 6, 1769–1820.
- [5] S. Brendle, *Rotationally symmetry of self-similar solutions to the Ricci flow* Invent. Math **194** (2013), no.3, 731–764.
- [6] E. Calabi, *On Ricci curvature and geodesics*, Duke Math. J. **34** (1967), 667–676.
- [7] H.-D. Cao, Q. Chen, *On locally conformally flat gradient steady Ricci solitons*, Trans. Amer. Math. Soc. **364** (2012), 2377–2391.
- [8] H.-D. Cao, H. Li, *A gap theorem for self-shrinkers of the mean curvature flow in arbitrary codimension*. Calc. Var. Partial Differential Equations. **46** (2013), 879–889.
- [9] J. Case, Y.-S. Shu, G. Wei, *Rigidity of quasi-Einstein metrics*, Differential Geom. Appl. **29** (2011), no. 1, 93–100.
- [10] G. Catino, C. Mantegazza, *The evolution of the Weyl tensor under the Ricci flow*. Ann. Inst. Fourier **61** (2011), no. 4, 1407–1435.
- [11] B.-L. Chen, *Strong uniqueness of the Ricci flow*, J. Differential Geom. **82** (2009), no. 2, 363–382.
- [12] X. Cheng, T. Mejia, and D. Zhou *Stability and compactness for complete f -minimal surfaces*. arXiv:1210.8076 (2012)
- [13] X. Cheng, D. Zhou, *Volume estimate about shrinkers*. Proc. Amer. Math. Soc. **141** (2013), 687–696.
- [14] T. Colding, W. Minicozzi, *Generic mean curvature flow I; generic singularities*. Ann. of Math. **175** (2012), 755–833.
- [15] Q. Ding, Y.L. Xin, *Volume growth, eigenvalue and compactness for self-shrinkers*. Asian J. Math. **17** (2012), 443–456
- [16] M. Eminenti, G. La Nave, C. Mantegazza, *Ricci Solitons: the equation point of view*, Manuscripta Math. **127** (2008), 345–367.
- [17] J. M. Espinar *Manifolds with density, applications and gradient Schrödinger operators*. arXiv:1209.6162v6. (2012)
- [18] E. M. Fan, *Topology of three-manifolds with positive P -scalar curvature*. Proc. Amer. Math. Soc. **136** (2008), no. 9, 3255–3261.
- [19] M. Fernández-López, E. García-Río, *A remark on compact Ricci solitons*, Math. Ann. **340** (2008), 893–896.
- [20] T. Frankel, *On the fundamental group of a compact minimal submanifold*, Ann. of Math. (2) **83** (1966), 68–73.
- [21] G. J. Galloway, *Compactness criteria for Riemannian manifolds*, Proc. Amer. Math. Soc. **84** (1982), no. 1, 106–110.
- [22] E. García-Río, D. Kupeli and B. Unal, *On a differential equation characterizing Euclidean spheres*, J. Differential Equations **194** (2003), 287–299.
- [23] R. E. Greene, H. Wu, *C^∞ approximations of convex, subharmonic, and plurisubharmonic functions*. Ann. Sci. École Norm. Sup. (4) **12** (1979), no.1, 47–84.
- [24] M. Gromov, *Isoperimetry of waists and concentration of maps*. Geom. Funct. Anal. **13** (2003), no. 1, 178–215.
- [25] R. S. Hamilton, *The Ricci flow on surfaces*, Mathematics and general relativity (Santa Cruz, CA, 1986), Contemp. Math., **71**. American Mathematical Society, Providence, 237–262 (1988).
- [26] G. Huisken, *Local and Global behaviour of hypersurfaces moving by mean curvature*, Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990), Proc. Sympos Pure Math. **54**, 175–191.
- [27] G. Huisken, *Asymptotic behaviour for singularities of the mean curvature flow*, J. Differential Geom. **31** (1990), no. 1, 285–299.

- [28] M. Kanai, *On a differential equation characterizing a Riemannian structure of a manifold*, Tokyo J. Math. **6** (1983), 143–151.
- [29] D.-S. Kim, Y.-H. Kim, *Compact Einstein warped product spaces with nonpositive scalar curvature*, Proc. Amer. Math. Soc. **131** (2003), no. 8, 2573–2576.
- [30] C. Li, *Index and topology of minimal hypersurfaces of \mathbb{R}^n* . Calc. Var. Partial Diff. Eq. (2017) **56**:180.
- [31] A. Naber, *Noncompact shrinking four solitons with nonnegative curvature*, J. Reine Angew. Math. **645** (2010), 125–153.
- [32] Z. Nehari, *Oscillation criteria for second-order linear differential equations*, Trans. Amer. Math. Soc. **85** (1957), 428–445.
- [33] M. Obata, *Certain conditions for a Riemannian manifold to be isometric with a sphere*, J. Math. Soc. Japan **14** (1962), 333–340.
- [34] G. Perelman, *Ricci flow with surgery on three manifolds*, arXiv:math.DG/0303109.
- [35] G. Perelman, *The entropy formula for the Ricci flow*, arXiv:math/0211159.
- [36] P. Petersen, W. Wylie. *Rigidity of gradient Ricci solitons*, Pacific J. Math. **241** (2009), no.2, 329–345.
- [37] P. Petersen, W. Wylie. *On the classification of gradient Ricci solitons*, Geom. Topol. **14** (2010), no. 4, 2277–2300.
- [38] A. Ros, *One-sided complete stable minimal surfaces*. J. Differential Geom. **74** (2006), no.1, 69-92.
- [39] A. Savo, *Index bounds for minimal hypersurfaces of the sphere*. Indiana Univ. Math. J. **59** (2010), no.3, 823-837.
- [40] R. Schoen, S.-T. Yau, *Harmonic Maps and the Topology of Stable Hypersurfaces and Manifolds with Non-negative Ricci Curvature*, Comm. Math. Helv. **51** (1976), 333–341.
- [41] R. Schoen, S.-T. Yau, *Compact group actions and the topology of manifolds with nonpositive curvature*, Topology **18** (1979), 361–380.
- [42] L. Shahriyari, *Translating graphs by mean curvature flow*, ProQuest LLC, Ann Arbor, MI, 2013, Thesis (Ph.D.)–The Johns Hopkins University.
- [43] K. Smoczyk, *A relation between mean curvature flow solitons and minimal submanifolds*, Math. Nachr. **229** (2001), 175–186.
- [44] Y. Tashiro, *Complete Riemannian manifolds and some vector fields*, Trans. Amer. Math. Soc. **117** (1965), 251–275.
- [45] G. Wei, W. Wylie, *Comparison geometry for the Bakry-Émery Ricci tensor*. J. Differential Geom. **83** (2009), no.2 377-405.
- [46] W. Wylie, *Complete shrinking Ricci solitons have finite fundamental group*, Proc. Amer. Math Soc. **136** (2008), no. 5, 1803–1806.
- [47] Qi S. Zhang *Extremal of log Sobolev inequality and W-entropy on noncompact manifolds* J. Funct. Anal. **263** (2012) no.7 2051–2101.
- [48] Z.-H. Zhang, *Gradient shrinking solitons with vanishing Weyl tensor*, Pacific J. Math. **242** (2009), no. 1, 189–200.
- [49] Z.-H. Zhang, *On the completeness of gradient Ricci solitons*, Proc. Amer. Math. Soc. **137** (2009), no. 8, 2755–2759.

Torino, May 23, 2019

.....